A protein related to the luminescence of the earthworm, Microscolex phosphoreus.

Ikuhiko Kin *1 and Yuichi Oba \dagger1

¹Chubu University (JAPAN) – Japan

Abstract

The earthworm *Microscolex phosphoreus* (Dugès, 1837), which was first described with specimens from Jardin des plantes de Montpellier in France, is one of the cosmopolitan luminous species [1]. This worm exudes yellow-greenish glowing mucus (λ max=527 nm) when stimulated mechanically or chemically. Its ecological function and mechanism of the luminescence remain unclear [2]. To clarify these questions, we established culturing method and examined the protein analysis. Protein mass spectrometry and transcriptomic analysis of the chloragogen tissue suggested that a 125 kDa protein is involved in the luminescence reaction.

Wampler, J. E. (1982). The bioluminescence system of *Microscolex phosphoreus* and its similarities to those of other bioluminescent earthworms (Oligochaeta). *Comparative Biochemistry and Physiology Part A: Physiology*, 71(4), 599-604.

Wampler, J. E., & Jamieson, B. G. (1986). Cell bound bioluminescence from *Pontodrilus* bermudensis, and its similarities to other earthworm bioluminescence. Comparative Biochemistry and Physiology Part A: Physiology, 84(1), 81-87.

Keywords: earthworm luciferase annelida

^{*}Speaker

[†]Corresponding author: yoba@isc.chubu.ac.jp